Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	3372	((544/122,331) or (514/235.8,252.19, 275)).CCLS.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2005/12/09 16:37








						OI ME	of Medicine Man				
PubMed Nucleo	otide Prot	ein Ge	enome Stru	cture	PopSe	et Ta	xonomy	OMIM Book			
Search PubMed	▼ for	· · · · · · · · · · · · · · · · · · ·	and the second s	* * * * * * * * * * * * * * * * * * *			Go	Clear			
	Limits	Pi	review/Index	History		Clip	oboard	Details			
		· · · · · · · · · · · · · · · · · · ·									
~	Display	Abstract	▼ Sort	<u> </u>	Save	Text	Clip Add	Order			
Entrez PubMed	☐ 1: Curr Opin Investig Drugs 2001 Sep;2(9):1213-9  Related Articles, Links										
	Lck inhibitors as a therapeutic approach to autoimmune disease and transplant rejection.										
PubMed Services	Kamens JS, Ratnofsky SE, Hirst GC.										
	Abbott Bioresearch Center, 100 Research Drive, Worcester, MA 01605 joanne.niewood@abbott.com										
Related Resources	T-cells play an important role in the pathogenesis of many diseases. These include diseases with large commercial markets and also with significant unmet medical needs, such as rheumatoid arthritis and asthma in addition to those with smaller markets such as organ transplantation, multiple sclerosis, inflammatory bowel diseases, type 1 diabetes, systemic lupus erythematosus and psoriasis. The use of currently available immunomodulatory agents is often limited by the appearance of dose-limiting side effects that result from the actions of these agents on non-lymphoid tissues. LSTRA cell kinase (lck), one of eight known members of the human src family of non-transmembrane protein tyrosine kinases, has a pivotal role in T-cell signaling. Lck expression is restricted to lymphoid cells, so an lck-selective inhibitor would be expected to have a significantly improved safety profile for the treatment of T-cell-driven diseases.  PMID: 11717807 [PubMed - indexed for MEDLINE]										
	Display	Abstract	▼ Sort		Save	Toyt	Clin Add	Order			


Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Jan 5. Prove 0.2 15 15 7 7 22341









**PubMed Services** 

Korade-Mirnics Z, Corey SJ.

Department of Pediatrics and Pharmacology, University of Pittsburgh School of Medicine, Pennsylvania, USA.

A concert of antigens, antibodies, cytokines, adhesion molecules, lipid factors, and their different receptors mediate leukocyte development and inflammatory responses. Regardless of the stimulus and receptor type, members of the Src family of protein tyrosine kinases (PTKs) play a critical role in initiating the numerous intracellular signaling pathways. Recruited and activated by the receptor, these Src PTKs amplify and diversify the signal. Multiple pathways arise, which affect cell migration, adhesion, phagocytosis, cell cycle, and cell survival. Essential nonredundant properties of Src PTKs have been identified through the use of gene targeting in mice or in the somatic cell line DT40. Because of their role in mediating leukocyte proliferation and activation, Src PTKs serve as excellent drug targets. Inhibitors of Src family members and dependent pathways may be useful in the treatment of human diseases similar to drugs known to inhibit other signal transduction pathways.

Related Resources

Publication Types:

- o Review
- Review, Tutorial

PMID: 11073097 [PubMed - indexed for MEDLINE]



Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer